

Topological charge and topological susceptibility in connection with translation and gauge invariance

A.M. Kotzinian ¹, O.Yu. Shevchenko ² and A.N. Sissakian ³

*Joint Institute for Nuclear Research
Dubna, Moscow region 141980, Russia*

Abstract

It is shown that the evaluation of the expectation value (EV) of topological charge density over θ -vacuum is reduced to investigation of the Chern-Simons term EV. An equation for this quantity is established and solved. EV of the topological charge density at an arbitrary θ occurs equal to zero and, as a consequence, topological susceptibility of both QCD and pure Yang-Mills vacua defined in a Wick sense is equal to zero, whereas when defined in a Dyson sense it differs from zero by the quantity proportional to the respective condensate of the chromomagnetic field. Thus, the usual Witten-Veneziano formula for the η' meson mass is modified.

¹E-mail address: Aram.Kotzinian@cern.ch

²E-mail address: shevch@nusun.jinr.ru

³E-mail address: sisakian@jinr.ru

The effects connected with the nontrivial topological configurations of the gauge fields attract a great attention in modern physics. In this respect the QCD topological susceptibility

$$\chi_{QCD} = \int d^4x \langle Tq(x)q(0) \rangle \quad (1)$$

is the quantity of a special importance because it enters as a key object in a lot of physical tasks, in particular, in such important puzzles as a famous $U(1)$ problem [1-5] (see [6] for a recent review) and the "spin crisis" [7]. In Eq.(1) $q(x)$ is the topological charge density

$$q(x) = \frac{g^2}{32\pi^2} \text{tr} F_{\mu\nu}^a(x) \tilde{F}_a^{\mu\nu}(x) \quad (2)$$

related with the Chern-Simons current, $K_\mu(x)$ by

$$q(x) = \partial^\mu K_\mu(x), \quad (3)$$

where

$$K_\mu = \frac{g^2}{32\pi^2} \epsilon^{\mu\nu\rho\sigma} A_\nu^a \left(F_{\rho\sigma}^a - \frac{g}{3} f_{abc} A_\rho^b A_\sigma^c \right). \quad (4)$$

It is well known, that topological susceptibility χ_{QCD} is equal to zero in all orders of perturbation theory and, also, that this quantity is just zero in the presence of even one massless quark (Crewther theorem [2]).

In this paper the consideration based on the fundamental translation and gauge symmetries will be performed which will allow to draw some unexpected conclusions about the topological charge and susceptibility.

Let us proof the following **statement**.

EV of the topological charge density $\langle \theta | q(0) | \theta \rangle = (1/VT) \langle \theta | Q | \theta \rangle$ over θ -vacuum with an arbitrary θ is equal to zero if EV of operator $K_\mu(0)$ over θ -vacuum exists, i.e.,

$$| \langle \theta | K_\mu(0) | \theta \rangle | < \infty, \quad (5)$$

where symbol $|\theta\rangle$ denotes the θ -vacuum state normalized to unity:

$$\langle \theta | \theta \rangle = 1. \quad (6)$$

This statement directly follows from translation invariance of θ -vacuum:

$$\begin{aligned} (\theta|q(0)|\theta) &= (\theta|\partial^\mu K_\mu(0)|\theta) \\ &= -i(\theta|[\hat{P}^\mu, K_\mu(0)]|\theta) = -i(P_\theta^\mu - P_\theta^\mu)(\theta|K_\mu(0)|\theta) = 0. \end{aligned} \quad (7)$$

The key point here is the condition (5) which, as we will see below, in $A_0 = 0$ gauge is equivalent to the condition

$$|(\theta|W_{CS}(0)|\theta)| < \infty, \quad (8)$$

where⁴

$$W_{CS}(t) \equiv \int d^3x K_0(x) = \frac{g^2}{32\pi^2} \int d^3x \left(\epsilon^{ijk} A_i^a \left(F_{jk}^a - \frac{g}{3} f_{abc} A_j^b A_k^c \right) \right) \quad (9)$$

is the Chern-Simons operator (see [8] for review). However, as we will see, within the conventional formulation of θ -vacuum theory rather amazing situation arises. On the one hand the condition (8) *is not satisfied* due to the gauge non-invariance of the operator W_{CS} with respect to the "large", topologically nontrivial gauge transformations. Nevertheless, despite EV $\langle \theta' | W_{CS}(0) | \theta \rangle$ is more singular function than $\delta(\theta' - \theta)$ at $\theta' \rightarrow \theta$ (namely, it behaves as $\delta'(\theta' - \theta)$ in this limit), the EV of the topological charge density over θ -vacuum is just zero again.

Since we deal with the gauge invariant quantity (EV of the topological charge) let us choose the Weil gauge

$$A_0 = 0, \quad (10)$$

which allows to essentially simplify a consideration. Choosing the periodic boundary conditions in the space directions (topology of a hypercylinder oriented along the time axis) one has

$$\int d^3x \partial^i K_i(t, \vec{x}) = 0, \quad (11)$$

and the expression for the topological charge

$$Q \equiv \int d^4x q(x) = \int d^4x \partial^\mu K_\mu \quad (12)$$

⁴ In the literature Chern-Simons term $W_{CS}[A]$ is also often called "collective coordinate" (see, for example, [5]) and is denoted by $X[A]$.

becomes (see, for example, [8,9])

$$Q = W_{CS}(t = \infty) - W_{CS}(t = -\infty).$$

EV $(\theta|\hat{O}|\theta)$ of an arbitrary operator $\hat{O}$ over θ -vacuum is defined as (see, for example [8,9])

$$(\theta|\hat{O}|\theta) = \frac{\langle\theta|\hat{O}|\theta\rangle}{\langle\theta|\theta\rangle} \quad (13)$$

where $|\theta\rangle$ is, simultaneously, the eigenfunction of the full QCD Hamiltonian H and of the unitary operator T_ν of the large gauge transformations with a winding number ν :

$$H|\theta\rangle = E_\theta|\theta\rangle, \quad (14)$$

$$T_\nu|\theta\rangle = e^{-i\theta\nu}|\theta\rangle, \quad (15)$$

i.e., the state $|\theta\rangle$ is, up to a phase multiplier, gauge invariant against the large gauge transformations. Notice also that on the contrary to (6) the states $|\theta\rangle$ are normalized as

$$\langle\theta'|\theta\rangle = \delta(\theta' - \theta). \quad (16)$$

so that the prescription (13) reads

$$\langle\theta'|\hat{O}|\theta\rangle = (\theta|\hat{O}|\theta)\delta(\theta' - \theta), \quad (17)$$

i.e., $(\theta|\hat{O}|\theta)$ is just the limit at $\theta' \rightarrow \theta$ of the multiplier at δ -function in the expression for $\langle\theta'|\hat{O}|\theta\rangle$.

Since we are interested in the quantity

$$(\theta|q(0)|\theta) = (VT)^{-1}(\theta|Q|\theta), \quad (18)$$

we will keep the normalization factor $(VT)^{-1}$. Using (11) and the Heisenberg equations one easily gets

$$\begin{aligned} (VT)^{-1}\langle\theta'|Q|\theta\rangle &= (VT)^{-1} \int dt e^{i(E_{\theta'} - E_\theta)t} \langle\theta'| \int d^3x q(0, \vec{x}) |\theta\rangle \\ &= 2\pi(VT)^{-1} \delta(E_{\theta'} - E_\theta) \langle\theta'| \dot{W}_{CS}(0) |\theta'\rangle \\ &= 2\pi(VT)^{-1} \delta(E_{\theta'} - E_\theta) \langle\theta'| -i[W_{CS}(0), H] |\theta\rangle \\ &= 2\pi i(VT)^{-1} \delta(E_{\theta'} - E_\theta) (E_{\theta'} - E_\theta) \langle\theta'| W_{CS}(0) |\theta\rangle, \end{aligned} \quad (19)$$

and, thus, the task now is to evaluate EV $\langle \theta' | W_{CS}(0) | \theta \rangle$.

The remarkable property of the Chern-Simons term is its transformation law under topologically nontrivial (often called "large" [8]) gauge transformations

$$A_i \rightarrow A_i^{\Omega_\nu} = \Omega_\nu A_i \Omega_\nu^{-1} + \partial_i \Omega_\nu \Omega_\nu^{-1} \quad (i = 1, 2, 3; \quad \Omega = \Omega(\vec{x})), \quad (20)$$

with topological index (winding number) ν . Namely, the quantity $W_{CS}[A]$ is not gauge invariant under (20) but transforms as

$$W_{CS}[A] \rightarrow W_{CS}[A^{\Omega_\nu}] = W_{CS}[A] + \nu, \quad (21)$$

i.e. it only shifts by the winding number ν of the respective gauge transformation.

The compatibility of the quantum

$$W_{CS}[A^{\Omega_\nu}] = T_\nu W_{CS}[A] T_\nu^+ = W_{CS}(A) + [T_\nu, W_{CS}(A)] T_\nu^{-1} \quad (22)$$

and classical (21) gauge transformation laws of the Chern-Simons term (9) gives rise to the commutation law ⁵

$$[T_\nu, W_{CS}(t)] = [T_\nu, W_{CS}(0)] = \nu T_\nu. \quad (23)$$

Now one already can evaluate $\langle \theta' | W_{CS}(0) | \theta \rangle$. Indeed, due to the unitarity of the operator T_ν and Eq. (15) one has

$$\langle \theta' | [T_\nu, W_{CS}(0)] | \theta \rangle = \left(e^{-i\nu\theta'} - e^{-i\nu\theta} \right) \langle \theta' | W_{CS}(0) | \theta \rangle. \quad (24)$$

On the other hand the commutation law (23) together with Eqs. (15), (16) give

$$\langle \theta' | [T_\nu, W_{CS}(0)] | \theta \rangle = \nu e^{-i\nu\theta} \delta(\theta - \theta'). \quad (25)$$

Comparing (24) and (25) one gets the basic for what follows equation

$$\frac{1}{\nu} \left(e^{-i\nu(\theta' - \theta)} - 1 \right) \langle \theta' | W_{CS}(0) | \theta \rangle = \delta(\theta' - \theta). \quad (26)$$

⁵Here one uses that $[T_\nu, H] = 0$.

So, one has to solve the equation

$$\frac{1}{\nu} \left(e^{-i\nu z} - 1 \right) f(z, \theta) = \delta_{2\pi}(z), \quad (27)$$

where $z \equiv \theta' - \theta$, $f(z, \theta) \equiv \langle \theta' | W_{CS}(0) | \theta \rangle$, and $\delta_{2\pi}(z) \equiv \delta(\theta' - \theta)$ is 2π -periodic δ -function. Expanding $f(z, \theta)$ and $\delta_{2\pi}(z)$ in the Fourier series

$$f(z, \theta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} \tilde{f}_n(\theta) e^{inz}, \quad \delta_{2\pi}(z) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} e^{inz}, \quad (28)$$

one easily obtains instead of (27) the difference equation for Fourier image $\tilde{f}$

$$\tilde{f}_{n+\nu}(\theta) - \tilde{f}_n(\theta) = \nu \quad (29)$$

with the solution

$$\tilde{f}_n(\theta) = n + C(\theta), \quad (30)$$

where $C(\theta)$ is some arbitrary function of θ .

Thus, the the general solution ⁶ of equation (25) reads ⁷

$$\langle \theta' | W_{CS}(0) | \theta \rangle = -i\delta'(\theta' - \theta) + C(\theta)\delta(\theta' - \theta), \quad (31)$$

where $\delta'(\theta' - \theta) = (i/2\pi) \sum n e^{inz}$.

Since

$$\lim_{\theta' \rightarrow \theta} \left(\frac{2\pi}{T} \delta(E_{\theta'} - E_{\theta}) \right) (E_{\theta'} - E_{\theta}) = (E_{\theta} - E_{\theta}) = 0, \quad (32)$$

the term $C(\theta)\delta(\theta' - \theta)$ in the solution (31) does not contribute to the coefficient at $\delta(\theta' - \theta)$ in the r.h.s. of (19) and, thus

$$(VT)^{-1} \langle \theta' | Q | \theta \rangle = 2\pi(VT)^{-1} \delta'(\theta' - \theta) [(E_{\theta'} - E_{\theta}) \delta(E_{\theta'} - E_{\theta})]. \quad (33)$$

⁶It is of importance that (31) is the solution of (26) only if the winding number ν is an arbitrary *integer* number (the existence of the fractional winding numbers was advocated in [2]). Otherwise, one can not perform the necessary change $n - \nu \rightarrow n$ in the sum $\sum n e^{i(n-\nu)z}$.

⁷Notice that for any *gauge invariant* operator $[T_{\nu}, O_{g.inv}] = 0$ and, therefore, Eq. (31) is replaced by $[\exp(-i\nu\theta') - \exp(-i\nu\theta)] \langle \theta' | O_{g.inv} | \theta \rangle = 0$ with the general solution $\langle \theta' | O_{g.inv} | \theta \rangle = C(\theta)\delta(\theta' - \theta)$.

Considering θ' as a variable whereas θ is kept fixed, one gets

$$(VT)^{-1}\langle\theta'|Q|\theta\rangle = -2\pi(VT)^{-1}\delta(\theta' - \theta)\frac{\partial E(\theta')}{\partial\theta'} \\ \times \left[\delta'(E_{\theta'} - E_{\theta})(E_{\theta'} - E_{\theta}) + \delta(E_{\theta'} - E_{\theta}) \right], \quad (34)$$

On the first sight the expression in the square brackets is equal to zero since usually $x\delta'(x) = -\delta(x)$. However, this is not correct conclusion and one has to properly work with the generalized function $(E_{\theta'} - E_{\theta})\delta'(E_{\theta'} - E_{\theta})$ when takes the limit $\theta' \rightarrow \theta$, i.e., $(E_{\theta'} - E_{\theta}) \rightarrow 0$. Indeed, let us consider the generalized function

$$\Delta(x) \equiv x\delta'(x). \quad (35)$$

Then

$$\begin{aligned} \Delta(0) &= \int_{-\infty}^{\infty} dx \delta(x) \Delta(x) \\ &\equiv \int_{-\infty}^{\infty} dx \delta(x) [x\delta'(x)] = - \int_{-\infty}^{\infty} dx \delta(x) [\delta(x) + x\delta'(x)] \\ &= - \int_{-\infty}^{\infty} dx \delta(x) \Delta(x) - \int_{-\infty}^{\infty} dx \delta(x) \delta(x) = -\Delta(0) - \delta(0), \end{aligned}$$

and, thus⁸

$$\Delta(0) \equiv [x\delta'(x)]|_{x=0} = -\frac{1}{2}\delta(0). \quad (36)$$

In particular,

$$\lim_{\theta' \rightarrow \theta} [(E_{\theta'} - E_{\theta})\delta'(E_{\theta'} - E_{\theta})] = -\frac{1}{2}\delta(E = 0) = -\frac{1}{2}\frac{T}{2\pi}. \quad (37)$$

In accordance with the general prescription (17) and Eqs. (18), (34), (37) one obtains

$$(\theta|q(0)|\theta) = -\frac{\partial E_{\theta}}{\partial\theta} \frac{2\pi}{VT} \left[-\frac{1}{2}\frac{T}{2\pi} + \frac{T}{2\pi} \right] = -\frac{1}{2}\frac{\partial\epsilon(\theta)}{\partial\theta}, \quad (38)$$

⁸The quantity $\Delta(x)$ is equal to $-\delta(x)$ as a generalized function only in the convolution with a function $F(x)$ satisfying the condition $xF'(x)|_{x=0} = 0$. However, this is just not the case for the choice $F(x) = \delta(x)$.

where

$$\epsilon(\theta) \equiv E_\theta/V \quad (39)$$

is the energy density of θ -vacuum.

On the other hand, it is well known that the energy density $\epsilon(\theta)$ is defined via the functional integral as (see, for example, [5,9])

$$\epsilon(\theta) = i(VT)^{-1} \ln W_\theta, \quad (40)$$

where

$$W(\theta) \equiv \int \mathcal{D}A \mathcal{D}[\bar{\psi}, \psi] \exp i [S_{QCD} + \theta Q]. \quad (41)$$

In this picture one has

$$(\theta|q(0)|\theta) = (VT)^{-1}(\theta|Q|\theta) = -i(VT)^{-1} \frac{\partial W[\theta]}{\partial \theta} W^{-1}[\theta] = -\frac{\partial \epsilon(\theta)}{\partial \theta}, \quad (42)$$

whereas the second derivative of $\epsilon(\theta)$ with respect to θ just produces the *topological susceptibility* - the connected part of the two-point correlator of the topological charge densities at zero momentum:

$$\chi_\theta = \int d^4x (\theta|Tq(x)q(0)|\theta)_{conn} = -(VT)^{-1} \frac{\partial^2 \ln W_\theta}{\partial \theta^2} = i \frac{\partial^2 \epsilon(\theta)}{\partial \theta^2}, \quad (43)$$

and

$$\chi_{QCD} = \chi_0. \quad (44)$$

It is easy to see now that the only way to reconcile Eqs. (38) and (42) is to put

$$(\theta|q(0)|\theta) = -\frac{\partial \epsilon(\theta)}{\partial \theta} = 0. \quad (45)$$

Then one can see that the topological susceptibility defined by (43) is also equal to zero

$$\chi_\theta = \chi_0 = 0. \quad (46)$$

Let us now attempt to realize obtained result. On the first glance Eq. (46) is in a severe contradiction with a standard point of view [2-6] that the quantity χ must differ from zero because it is directly connected with the solution of $U(1)$ -problem and mass of η' -meson is explicitly expressed via topological susceptibility. However, the situation perhaps is not so bad. The point is that there exist *two different topological susceptibilities* in accordance with the different sense of the time-ordering operation in the respective correlators.

Let us remind that the functional integral representation (40)-(42) for a correlator means that the respective T-product in this correlator must be realized as the Wick (T_W) time-ordering operation, in which all the derivatives are applied after the calculation of the field convolutions (see [5] for the excellent review on this question). On the contrary, the Dyson T-ordering of two arbitrary operators $A(x)$ and $B(y)$ (does not matter composite or not) simply looks as

$$T_D[A(x)B(y)] = \theta(x_0 - y_0)A(x)B(y) + \theta(y_0 - x_0)B(y)A(x). \quad (47)$$

So, actually Eq. (46) have to be read as

$$\chi_\theta^W = \chi_0^W = 0, \quad (48)$$

where

$$\chi_\theta^W \equiv \int d^4x [(\theta|T_W[q(x)q(0)]|\theta)_{conn}], \quad (49)$$

but *it does not mean at all* that the Dyson topological susceptibility

$$\chi_\theta^D = \int d^4x [(\theta|T_D q(x)q(0)|\theta) - (\theta|q(0)|\theta)^2] = \int d^4x (\theta|T_D q(x)q(0)|\theta)_{conn} \quad (50)$$

is also equal to zero.

Indeed, the connection between Wick and Dyson susceptibilities was found in [5] using the stationary perturbation theory in powers of θ with a result

$$\chi_0^W = i \frac{\partial^2 \epsilon}{\partial \theta^2} \Big|_{\theta=0} = \chi_0^D + i \langle 0 | \left(\frac{g^2}{8\pi^2} \vec{B}_a \right)^2 | 0 \rangle, \quad (51)$$

i.e., these susceptibilities differ by the condensate of the chromomagnetic field $\vec{B}_a$. Thus, even despite that in accordance with (48) the quantity χ_0^W

equal to zero, the Dyson topological susceptibility χ_D^0 can differ from zero by the nonzero value of the chromomagnetic condensate:

$$\chi_0^D = -i \langle 0 | \left(\frac{g^2}{8\pi^2} \vec{B}_a \right)^2 | 0 \rangle. \quad (52)$$

On the other hand it is well known that namely Dyson T-product (not Wick!) can be represented as the sum over intermediate states with a result

$$\begin{aligned} \int d^4x e^{ik \cdot x} \langle 0 | T_D(A(x)B(0)) | 0 \rangle_{conn} &= \sum_n \langle 0 | A(0) | n(\vec{k}) \rangle \langle n(\vec{k}) | B(0) | 0 \rangle \frac{i}{k^2 - m_n^2} \\ &+ \text{two particle contributions} \quad (k^2 = k_0^2 - \vec{k}^2). \end{aligned} \quad (53)$$

As it was shown by Witten [3] (see [11], sect 5.1, for review) only the one particle contributions survive in the sum over intermediate states in the large N_c limit. So,

$$\begin{aligned} \chi_0^D |_{QCD} &= \int d^4x \langle 0 | T_D q(x) q(0) | 0 \rangle_{conn} |_{QCD} \\ &= \lim_{k \rightarrow 0} \left[\sum_{n=messons} |\langle 0 | q(0) | n(\vec{k}) \rangle|^2 \frac{i}{k^2 - m^2} + \sum_{l=glubols} |\langle 0 | q(0) | l(\vec{k}) \rangle|^2 \frac{i}{k^2 - m^2} \right] \\ &= \chi_0^D |_{YM} + \sum_{n=messons} |\langle 0 | q(0) | n \rangle|^2 \frac{-i}{m_n^2}, \end{aligned} \quad (54)$$

and, in the leading order of χ_{PT} where [6]

$$\langle 0 | q(0) | \eta' \rangle = \langle 0 | \frac{1}{2N_f} \partial^\mu J_\mu^5 | \eta' \rangle = \frac{1}{2N_f} f_{\eta'} m_{\eta'}^2 \quad (55)$$

one gets instead of Witten-Veneziano [3,4] formula

$$\begin{aligned} m_{\eta'}^2 &= -i \frac{4N_f^2}{f_{\eta'}^2} (\chi_0^D |_{YM} - \chi_0^D |_{QCD}) \\ &= \frac{4N_f^2}{f_{\eta'}^2} \left[\langle 0 | \left(\frac{g^2}{8\pi^2} \vec{B}_a \right)^2 | 0 \rangle |_{QCD} - \langle 0 | \left(\frac{g^2}{8\pi^2} \vec{B}_a \right)^2 | 0 \rangle |_{YM} \right]. \end{aligned} \quad (56)$$

It is of importance and seems to be a serious argument in support of our model independent consideration (based on the general principles of translational and gauge invariance) that very similar results concerning η' -meson

mass was obtained within the different QCD inspired models. These are Cheshire cat principle model [12] and, also, squeezed gluon vacuum [13] and monopole vacuum [14] models (compare⁹ Eq. (56) with Eq. (14) of ref. [12] and, especially, with Eqs. (18) and (326) of refs. [13] and [14], respectively).

Thus, we get rather unexpected result: Wick topological susceptibility is equal to zero whereas the Dyson topological susceptibility just proportional to the chromomagnetic condensate. The last circumstance allows one to get the mass formula (56) for the η' -meson which directly express its mass via the difference of the respective chromomagnetic condensates with and without quark inclusion.

Acknowledgment

The authors are grateful to E. Kuraev, S. Nedelko, V. Pervushin, I. Solovtsov, O. Teryaev and G. Veneziano for fruitful discussions.

References

- [1] G. 't Hooft, Phys. Rep. **142** (1986) 357.
- [2] R.J. Crewther, Phys. Lett. **B70** (1977) 349; Acta Phys. Austr. Suppl. **19** (1978) 47; Riv. Nuovo. Cim. **2N8** (1979) 63.
- [3] E. Witten, Nucl. Phys. **B156** (79) 269.
- [4] G. Veneziano, Nucl. Phys. **B159** (79) 213
- [5] D.I. Dyakonov and M.I. Eides, Sov. Phys. JETP **54(2)** (1981) 232
- [6] B.L. Ioffe, Phys. Atom. Nucl. **62** (1999) 2025
- [7] S. Narison, G.M. Shore, G. Veneziano, Nucl. Phys. **B433** (1995) 209
- [8] R. Jackiw in "Current algebra and anomalies", eds S. Trieman, R. Jackiw, B. Zumino and E. Witten (World Scientific, Singapore, 1985)

⁹Comparing these formulas one has to use that $\alpha_s = g^2/4\pi$ and $f_{\eta'} = \sqrt{2N_f}f_\pi$.

- [9] R. Rajaraman, "An Introduction to Solitons and Instantons in Quantum Field Theory", North-Holland, 1982;
F.J. Yndurain, "Quantum Chromodynamics. An Introduction to the Theory of Quarks and Gluons", Springer-Verlag, 1983 (section 38).
- [10] K. Huang, "Quarks, Leptons and Gauge Fields", World Scientific, Singapore, 1982
- [11] G.A. Christos, Phys. Rep. **116** (1984) 251.
- [12] H.B. Nielsen et al., Phys.Lett. **B281** (1992) 345.
- [13] D. Blaschke et al., Phys.Lett. **B397** (1997) 129.
- [14] V.N. Pervushin, "Dirac variables in gauge theories", hep-th/0109218.